UNIT 12 : LINEAR HOMOGENEOUS DIFFERENTIAL EQUATION OF HIGHER ORDER

UNIT STUCTURE

12.1 Learning Objectives

12.2 Introduction

12.3 Linear Homogeneous Equation

12.3.1 Equation Reducible to Homogeneous form

12.4 Method of Variation of Parameters

12.5 Let Us Sum Up

12.6 Further Readings

12.7 Answers To Check Your Progress

12.8 Model Questions

12.1 LEARNING OBJECTIVES

After going through this unit, you will able to know

- linear homogeneous equation
- equation reducible to homogeneous form
- method of variation of parameters.

12.2 INTRODUCTION

In unit 11, we have discussed about solution of linear differential equation of higher order with constant coefficients. In this unit, we will discuss solution of linear homogeneous differential equations. We shall also discuss the method of variation of parametres for solving linear differential equation with constant coefficients.

12.3 LINEAR HOMOGENEOUS EQUATION

A differential equation of the form

$$a_n x^n \frac{d^n y}{dx^n} + a_{n-1} x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_2 x^2 \frac{d^2 y}{dx^2} + a_1 x \frac{dy}{dx} + a_0 y = X \quad \text{... (1)}$$

where $a_0, a_1, a_2, \cdots, a_{n-1}, a_n$ are constants and X is a function of x is called a homogeneous linear equation.

The equation is reducible to a linear equation with constants coefficients with the help of substitution $x = e^z$ which can be solved the methods described in the section 7.3.

Let $x = e^z$ i.e., $z = \log x$, then

$\frac{dy}{dx} = \frac{dy}{dz} \cdot \frac{dz}{dx} = \frac{1}{x} \frac{dy}{dz}$ so that $x \frac{dy}{dx} = \frac{dy}{dz} = Dy$ where $D = \frac{d}{dz}$

$$\frac{d^2 y}{dx^2} = \frac{d}{dx}\left(\frac{1}{x}\frac{dy}{dz}\right)$$

$$= -\frac{1}{x^2}\frac{dy}{dz} + \frac{1}{x}\frac{d^2 y}{dz^2}\frac{dz}{dx}$$

$$= -\frac{1}{x^2}\frac{dy}{dz} + \frac{1}{x^2}\frac{d^2 y}{dz^2}$$

or $x^2 \frac{d^2 y}{dx^2} = -\frac{dy}{dz} + \frac{d^2 y}{dz^2} = \left(D^2 - D\right)y = D(D-1)y$

Similarly, $x^3 \frac{d^3 y}{dx^3} = \left(D^2 - D\right)y = D(D-1)(D-2)y$

and in general $x^n \frac{d^n y}{dx^n} = D(D-1)(D-2)\cdots(D-n+1)y$

Subtituting for $x\frac{dy}{dx}, x^2 \frac{d^2 y}{dx^2}$,... the equation (1) reduces to a linear equation with constants coefficients which is solvable by the methods already discussed.

Example 1 : Solve $x^2 \frac{d^2 y}{dx^2} - x\frac{dy}{dx} + y = \log x$

Solution : Let $x = e^z$ i.e., $z = \log x$

$\frac{dy}{dx} = \frac{dy}{dz} \cdot \frac{dz}{dx} = \frac{1}{x} \frac{dy}{dz}$ so that $x \frac{dy}{dx} = \frac{dy}{dz} = Dy$ where $D = \frac{d}{dz}$

so, $x^2\frac{d^2y}{dx^2} = -\frac{dy}{dz} + \frac{d^2y}{dz^2} = \left(D^2 - D\right)y = D(D-1)y$

The given equation reduces

$$\left(D(D-1) - D + 1\right)y = z$$

or $(D-1)^2 = y$

C.F $= (Az + B)e^z = (A\log x + B)x$

P.I $= \frac{1}{(D-1)^2}z$

$= (1-D)^{-2}z$

$= (1+2D)z$

$= z + 2$

$= \log x + 2$

The general solution is $y = (A\log x + B)x + \log x + 2$.

Example 2 : Solve $x^3\frac{d^3y}{dx^3} + 3x^2\frac{d^2y}{dx^2} + x\frac{dy}{dx} = x^3\log x$

Solution : Let $x = e^z$ i.e., $z = \log x$

$\frac{dy}{dx} = \frac{dy}{dz}.\frac{dz}{dx} = \frac{1}{x}\frac{dy}{dz}$ so that $x\frac{dy}{dx} = \frac{dy}{dz} = Dy$ where $D = \frac{d}{dz}$

so, $x^2\frac{d^2y}{dx^2} = -\frac{dy}{dz} + \frac{d^2y}{dz^2} = \left(D^2 - D\right)y = D(D-1)y$

The given equation reduces

$\left[D(D-1)(D-2) + 3D(D-1) + D)\right]y = e^{3z}z$

or $D^3y = e^{3z}z$

C.F $= \left(A + Bz + Cz^2\right)e^{0.z} = A + B\log x + C(\log x)$

P.I $= \frac{1}{D^3}\left(e^{3z}z\right)$

$= e^{3z}\frac{1}{(D+3)^3}z$

$= e^{3z}\left(1 + \frac{D}{3}\right)^{-3}z$

$= \frac{e^{3z}}{27}(1-D)z$

$$= \frac{e^{3z}}{27}(z-1)$$

$$= \frac{x^3}{27}(\log x - 1)$$

The general solution is $y = A + B\log x + C(\log x)^2 + \frac{x^3}{27}(\log x - 1)$.

12.3.1 Equations Reducible to Homogeneous form

An equation of the form

$$(a+bx)^n \frac{d^n y}{dx^n} + P_1(a+bx)^{n-1}\frac{d^{n-1}y}{dx^{n-1}} + \cdots + P_{n-1}(a+bx)\frac{dy}{dx} + P_n y = X$$

where $P_1, P_2, \cdots, P_n$ are all constants,can be reduced to the homogeneous linear form by putting $a+bx = e^z$ and can be solved by the method discussed in 7.4.

Example 3 : Solve $(1+x)^2\frac{d^2y}{dx^2} + (1+x)\frac{dy}{dx} + y = 2\sin\log(1+x)$

Solution : Let $1+x = e^z$ i.e., $z = \log(1+x)$

$\frac{dy}{dx} = \frac{dy}{dz}.\frac{1}{1+x}$ or $(1+x)\frac{dy}{dx} = \frac{dy}{dz} = Dy$

similarly, $(1+x)^2\frac{d^2y}{dx^2} = D(D-1)y$

The given equation becomes

$$D(D-1)y + Dy + y = 2\sin z$$

or $(D^2+1)y = 2\sin z$

Auxiliary equation is $m^2 + 1 = 0$

or $m = \pm i$

C.F $= A\cos z + B\sin z = A\cos\log(1+x) + B\sin\log(1+x)$

P.I $= \frac{1`}{D^2+1}(2\sin z)$

$$= 2\mathrm{Im}\left(\frac{1`}{D^2+1}e^{iz}\right)$$

$$= 2\mathrm{Im}\left(e^{iz}\frac{1`}{(D+i)^2+1}1\right)$$

$$= 2\mathrm{Im}\left(e^{iz} \frac{1`}{D^2 + 2iD} 1 \right)$$

$$= 2\mathrm{Im}\left(e^{iz} \frac{1}{2iD}\left(1 + \frac{D}{2i}\right)^{-1} 1 \right)$$

$$= 2\mathrm{Im}\left(e^{iz} \frac{1}{2iD} 1 \right)$$

$$= 2\mathrm{Im}\left(e^{iz} \frac{z}{2i} \right)$$

$$= \mathrm{Im}\left(\frac{z\cos z + 2i\sin z}{i} \right)$$

$$= -z\cos z$$

$$= -\log(1+x)\cos\log(1+x)$$

The general solution is

$$y = A\cos\log(1+x) + B\sin\log(1+x) - \log(1+x)\cos\log(1+x).$$



CHECK YOUR PROGRESS - 1

Q 1 : Solve the following differential equations :

(a) $x^3 \frac{d^3y}{dx^3} + 2x^2 \frac{d^2y}{dx^2} + 2y = 10\left(x + \frac{1}{x}\right)$

(b) $\left(x^2D^2 - 3xD + 4\right)y = 2x^2$

(c) $\left(x^2D^2 - xD + 4\right)y = \cos(\log x) + x\sin(\log x)$

(d) $(x+a)^2 \frac{d^2y}{dx^2} - 4(x+a)\frac{dy}{dx} + 6y = x$

12.4 METHOD OF VARIATION OF PARAMETERS

Theorem : For the equation $y'' + Py' + Qy = X$ where P, Q and X being functions of x, the P.I is given by $-y_1\int \frac{y_2X}{W}dx + y_2\int \frac{y_1X}{W}dx$. Here y_1 and y_2 are solutions of $y'' + Py' + Qy = 0$ and $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$, the Wronskian of y_1, y_2 named after the Polish mathematician and philospher Hoene Wronsky.



Hoene Wronsky (1776-1853) was a Polish mathematician and philospher

Proof : Let the complementary function of $y'' + Py' + Qy = X$ be $y = c_1 y_1 + c_2 y_2$. Here c_1 and c_2 can be regarded as parametric functions $u(x)$, $v(x)$.

Let the P.I be $y = u(x)y_1 + v(x)y_2$

Differentiating w.r.t x, we get

$y' = uy_1' + u'y_1 + vy_2' + v'y_2$

Choose u & v such that $u'y_1 + v'y_2 = 0$ so that

$$y' = uy_1' + vy_2'$$

and $y'' = u'y_1' + uy_1'' + v'y_2' + vy_2''$

Since y_1, y_2 are roots of $y'' + Py' + Qy = 0$, we have

$$y_1'' + Py_1' + Qy_1 = 0$$

and $$y_2'' + Py_2' + Qy_2 = 0$$

The given equation becomes

$u'y_1' + uy_1'' + v'y_2' + vy_2'' + P(uy_1' + vy_2') + Qy = X$

or $u(y_1'' + Py_1' + Qy_1) + v(y_2'' + Py_2' + Qy_2)$

$$-Q(uy_1 + vy_2) + u'y_1' + v'y_2' + Qy = X$$

or $u'y_1' + v'y_2' - Q(uy_1 + vy_2) + Qy = X$

or $u'y_1' + v'y_2' - Qy + Qy = X \quad (y = uy_1 + vy_2)$

or $u'y_1' + v'y_2' - X = 0$ ... (1)

Also, $u'y_1 + v'y_2 + 0.X = 0$ ... (2)

Solving (1) and (2), we get

$$\frac{u'}{y_2} = \frac{v'}{-y_1} = \frac{X}{y_1'y_2 - y_1y_2'} = \frac{X}{-W}$$

Hence $u' = \frac{X}{-W}y_2$ and $v' = \frac{X}{W}y_1$

i.e., $u = -\int \frac{y_2X}{W}dx$ and $v = \int \frac{y_1X}{W}dx$

and P. I $= -y_1\int \frac{y_2X}{W}dx + y_2\int \frac{y_1X}{W}dx$

Hence proved.

Example 4 : Solve $\frac{d^2y}{dx^2} + 4y = \tan 2x$

Solution : Auxiliary equation is $m^2 + 4 = 0$

or $m = \pm 2i$

C.F $= A\cos 2x + B\sin 2x$

$= Ay_1 + By_2$ where $y_1 = \cos 2x$ and $y_2 = \sin 2x$ and $X = \tan 2x$.

Wronskian, $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} \cos 2x & \sin 2x \\ -2\sin 2x & 2\cos 2x \end{vmatrix} = 2$.

P. I $= -y_1 \int \frac{y_2 X}{W} dx + y_2 \int \frac{y_1 X}{W} dx$

$= -\cos 2x \int \frac{\sin 2x \tan 2x}{2} dx + \sin 2x \int \frac{\cos 2x \tan 2x}{2} dx$

$= -\frac{\cos 2x}{2} \int \frac{1 - \cos^2 2x}{\cos 2x} dx + \sin 2x \int \frac{\sin 2x}{2} dx$

$= -\frac{1}{2}\cos 2x \int (\sec 2x - \cos 2x) dx - \frac{1}{4}\sin 2x \cos 2x$

$= -\frac{1}{2}\cos 2x \left\{ \frac{1}{2}\log(\sec 2x + \tan 2x) - \frac{1}{2}\sin 2x \right\} - \frac{1}{4}\sin 2x \cos 2x$

$= -\frac{1}{4}\cos 2x \left\{ \frac{1}{2}\log(\sec 2x + \tan 2x) - \sin 2x \right\} - \frac{1}{4}\sin 2x \cos 2x$

$= -\frac{1}{4}\cos 2x \log(\sec 2x + \tan 2x)$

The general solution is

$y = A\cos 2x + B\sin 2x - \frac{1}{4}\cos 2x \log(\sec 2x + \tan 2x)$.

Example 5 : Solve $\frac{d^2 y}{dx^2} + a^2 y = \sec ax$

Solution: Auxiliary equation is $m^2 + a^2 = 0$

or $m = \pm ia$

C.F $= A\cos ax + B\sin ax$

Let $y_1 = \cos ax$ and $y_2 = \sin ax$

Wronskian, $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} \cos ax & \sin ax \\ -a\sin ax & a\cos ax \end{vmatrix} = a$ and $X = \sec ax$.

P. I $= -y_1 \int \frac{y_2 \sec ax}{a} dx + y_2 \int \frac{y_1 \sec ax}{a} dx$

$$= -\frac{1}{2}\cos ax \int \frac{\sin ax}{\cos ax} dx + \frac{\sin ax}{a} \int dx$$

$$= \frac{1}{a^2}\cos ax \log \cos ax + \frac{x}{a}\sin ax$$

The general solution is

$$y = A\cos ax + B\sin ax + \frac{1}{a^2}\cos ax \log \cos ax + \frac{x}{a}\sin ax .$$

Example 6 : Solve by the method of variation of parameters :

$$y'' - 2y' + y = e^x \log x$$

Solution : The given equation can be written as

$$\left(D^2 - 2D + 1\right)y = e^x \log x$$

The auxiliary equation is $m^2 - 2m + 1 = 0$

or $(m-1)^2 = 0$

or $m = 1,1$

C. F $= (Ax + B)e^x$

Let $y_1 = e^x, y_2 = xe^x, X = e^x$ and $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} e^x & xe^x \\ e^x & e^x + xe^x \end{vmatrix}$

$$= e^{2x}\begin{vmatrix} 1 & x \\ 1 & 1+x \end{vmatrix} = e^{2x}$$

P. I $= -y_1 \int \frac{y_2 X}{W} dx + y_2 \int \frac{y_1 X}{W} dx$

$$= -e^x \int \frac{xe^{2x}\log x}{e^{2x}} dx + xe^x \int \frac{e^x . e^x \log x}{e^{2x}} dx$$

$$= -e^x \int x \log x dx + xe^x \int \log x dx$$

$$= -e^x \left\{ \log x . \frac{x^2}{2} - \int \frac{1}{x} . \frac{x^2}{2} dx \right\} + xe^x \left\{ x \log x - \int dx \right\}$$

$$= -e^x \left\{ \frac{x^2}{2}\log x - \frac{x^2}{4} \right\} + xe^x (\log x - 1)$$

$$= \frac{1}{4}x^2 e^x \{2\log x + 1 + 4(\log x - 1)\}$$

$$= \frac{1}{4}x^2 e^x (2\log x - 3)$$

The general solution is

$y = (Ax + B)e^x + \frac{1}{4}x^2 e^x (2\log x - 3)$.

Example 7 : Solve by method of varation of parameters :

$$\frac{d^2y}{dx^2} - y = \frac{2}{1+e^x}$$

Solution : Given, $\frac{d^2y}{dx^2} - y = \frac{2}{1+e^x}$

A. E is $m^2 - 1 = 0$

or $m^2 = 1$

or $m = \pm 1$

C.F $= C_1 e^x + C_2 e^{-x}$

and $y^1.y_2' - y_1'.y_2 = -e^x.e^{-x} - e^x.e^{-x} = -2$

$$u = \int \frac{-y^2 X}{y_1.y_2' - y_1'.y_2} dx = -\int \frac{e^{-x}}{-2} \times \frac{2}{1+e^x} dx$$

$$= \int \frac{e^{-x}}{1+e^x} dx$$

$$= \int \frac{dx}{e^x(1+e^x)}$$

$$= \int \left(\frac{1}{e^x} - \frac{1}{1+e^x}\right) dx$$

$$= \int e^{-x} dx - \int \frac{e^{-x}}{e^{-x}+1} dx$$

$$= -e^{-x} + \log(e^{-x}+1)$$

$$v = \int \frac{y_1 X}{y_1.y_2' - y_1'.y_2} dx = \int \frac{e^{-x}}{-2} \times \frac{2}{1+e^x} dx$$

$$= \int \frac{e^x}{e^x+1} dx$$

$$= -\log(e^x+1)$$

P.I $= u.y_1 + v.y_2$

$= \left[-e^{-x} + \log(e^{-x}+1)\right]e^x - e^{-x}\log(1+e^x)$

$= -1 + e^x \log(e^{-x}+1) - e^{-x}\log(1+e^x)$

The general solution is

$y = C_1 e^x + C_2 e^{-x} - 1 + e^x \log\left(e^{-x}+1\right) - e^{-x}\log\left(1+e^x\right).$



CHECK YOUR PROGRESS - 2

Q 1 : Apply the method of variation of parameters to solve.

(i) $y_2 + a^2 y = \cot ax$

(ii) $y_2 - 2y_1 + y = xe^x \log x, x > 0$

(iii) $\left(D^2 - 2D + 2\right)y = e^x \tan x$

(iv) $y^{//} + 3y^{/} + 2y = x + \cos x$



12.5 LET US SUM UP

- A differential equation of the form

$$a_n x^n \frac{d^n y}{dx^n} + a_{n-1} x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_2 x^2 \frac{d^2 y}{dx^2} + a_1 x \frac{dy}{dx} + a_0 y = X$$

...(1)

where $a_0, a_1, a_2, \cdots, a_{n-1}, a_n$ are constants and X is a function of x is called a homogeneous linear equation.

The equation is reducible to a linear equation with constants coefficients with the help of substitution $x = e^z$ which can be solved the methods described in the section 12.3.

Let $x = e^z$ i.e., $z = \log x$

$$x\frac{dy}{dx} = \frac{dy}{dz} = Dy$$

$$x^2 \frac{d^2 y}{dx^2} = -\frac{dy}{dz} + \frac{d^2 y}{dz^2} = \left(D^2 - D\right)y = D(D-1)y$$

Similarly, $x^3 \frac{d^3 y}{dx^3} = \left(D^2 - D\right)y = D(D-1)(D-2)y$

and in general $x^n \frac{d^n y}{dx^n} = D(D-1)(D-2)\cdots(D-n+1)y$

The substitution of these values in the given equation reduces the given homogeneous equation to a differential equation with constants coefficients.

- Method of variation of parameter to find particular integral of

$P\frac{d^2y}{dx^2}+Q\frac{dy}{dx}+Ry=X$ is as follows :

Let complementary function(C.F) $= Ay_1 + By_2$

then particular integral(P.I) $= uy_1 + vy_2$

where $u=\int\frac{-y_2X}{W}dx$ and $v=\int\frac{y_1X}{W}dx$, $W=\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$.

General solution =C.F +P.I.



12.6 FURTHER READINGS

1) Tenenbaum, Morris and Pollard, Harry. Ordinary Differential Equations, Dover Publications, Inc, N.Y 11501.

2) Differential Equations: An Introduction with Mathematica. Clay C.Ross, second Edition, Springer.

3) Differential Equations: Linear, Nonlinear, ordinary, partial. A.C. King, J.Billinggham and S. R. Otto, Cambridge-2003.

4) Differential equations with Application and Programs. S. Balachandra Rao and H.R. Anuradha, University Press (India) Limited 1996.



12.7 ANSWERS TO CHECK YOUR PROGRESS

CHECK YOUR PROGRESS – 1

Ans to Q No 1 : (b) Given that $\left(x^2D^2-3xD+4\right)y=2x^2$ ... (1)

Let $x=e^z$ or $z=\log x$ and $D_1=\frac{d}{dz}$.

Then (1) becomes

$\{D_1(D_1-1)-3D_1+4\}y = 2e^{2z}$

or $(D_1-2)^2 y = 2e^{2z}$

Auxiliary equation is $(m-2)^2 = 0$

or $m = 2,2$

C. F $= (C_1 + C_2 z)e^{2z} = (C_1 + C_2 z)(e^z)^2 = (C_1 + C_2 \log x)x^2$

P. I $= \dfrac{1}{(D_1-2)^2} 2e^{2z}$

$= 2\dfrac{z^2}{2!}e^{2z} = z^2(e^z)^2$

$= (\log x)^2 x^2$

Complete solution is

$y = (C_1 + C_2 \log x)x^2 + (\log x)^2 x^2$.

(c) Putting $x = e^z$ and denoting $\dfrac{d}{dz}$ by D_1, the given equation becomes

$[D_1(D_1-1) - D_1 + 4]y = \cos z + e^z \sin z$

or $[D_1^{\,2} - 2D_1 + 4]y = \cos z + e^z \sin z$

Auxiliary equation is $m^2 - 2m + 4 = 0$

or $m = 1 \pm \sqrt{3}i$

C.F $= e^z[C_1 \cos \sqrt{3}z + C_2 \sin \sqrt{3}z]$

$= x[C_1 \cos \sqrt{3}(\log x) + C_2 \sin \sqrt{3}(\log x)]$

P.I $= \dfrac{1}{D_1^{\,2} - 2D_1 + 4}\cos z + \dfrac{1}{D_1^{\,2} - 2D_1 + 4} e^z \sin z$

$= \dfrac{1}{-1^2 - 2D_1 + 4}\cos z + e^z \dfrac{1}{(D_1+1)^2 - 2(D_1+1) + 4}\sin z$

$= \dfrac{1}{3 - 2D_1}\cos z + e^z \dfrac{1}{D_1^{\,2} + 4}\sin z$

$= \dfrac{(3 + 2D_1)}{9 - 4D_1^{\,2}}\cos z + e^z \dfrac{1}{-1^2 + 3}\sin z$

$= \dfrac{3\cos z - 2\sin z}{9+4} + \dfrac{e^z \sin z}{2}$

$$= \frac{1}{13}\left[3\cos(\log x) - 2\sin(\log x)\right] + \frac{x}{2}\sin(\log x)$$

Hence the complete solution is

$$y = x\left[C_1\cos\sqrt{3}(\log x) + C_2\sin\sqrt{3}(\log x)\right] + \frac{1}{13}\left[3\cos(\log x) - 2\sin(\log x)\right] + \frac{x}{2}\sin(\log x)$$

(d) Putting $x + a = e^z$ and denoting $\frac{d}{dx}$ by D ,the equation becomes

$\left[D(D-1) - 4D + 6\right]y = e^z - a$

or $\left[D^2 - 5D + 6\right]y = e^z - a$

Auxiliary equation is $m^2 - 5m + 6 = 0$

or $m = 3,2$

C.F $= C_1e^{3z} + C_2e^{2z} = C_1(x+a)^3 + C_2(x+a)^2$

P.I $= \frac{1}{D^2 - 5D + 6}.e^z - \frac{1}{D^2 - 5D + 6}.a$

$= \frac{1}{(D-3)(D-2)}.e^z - \frac{1}{(D-3)(D-2)}.a$

$= \frac{e^z}{(1-3)(1-2)} - a\frac{1}{(D-3)(D-2)}e^{0z}$

$= \frac{1}{3}e^z - a\frac{1}{(0-3)(0-2)}e^{0z}$

$= \frac{1}{3}e^z - a\frac{1}{6}e^{0z}$

$= \frac{1}{3}e^z - \frac{1}{6}a$

$= \frac{1}{3}(x+a) - \frac{1}{6}a$

Hence, the complete solution is

$$y = C_1(x+a)^3 + C_2(x+a)^2 + \frac{1}{3}(x+a) - \frac{1}{6}a$$

CHECK YOUR PROGRESS – 2

Ans to Q No 1 : (i) Given, $y_2 + a^2y = \cot ax$ ... (1)

Auxiliary equation is $m^2 + a^2 = 0$

or $m = \pm ia$

C.F $= C_1\cos ax + C_2\sin ax$

Here $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} \cos ax & \sin ax \\ -a\sin ax & a\cos ax \end{vmatrix} = a \neq 0$

Then P. I $= u.y_1 + v.y_2$

where $u = -\int \frac{y_2.X}{W} dx = -\int \frac{\sin ax \cot ax}{a} dx = -\frac{1}{a}\int \cos ax dx = -\frac{\sin ax}{a^2}$

and $v = \int \frac{y_1.X}{W} dx = \int \frac{\cos ax \cot ax}{a} dx = \frac{1}{a}\int \frac{1-\sin^2 ax}{\sin ax} dx$

$$= \frac{1}{a}\int (\cos ecax - \sin ax) dx$$

$$= \frac{1}{a}\int (\cos ecax - \sin ax) dx = \frac{1}{a^2}\left(\log \tan \frac{ax}{2} + \cos ax\right)$$

P. I $= \cos ax \times \left(-\frac{1}{a^2}\right) \times \frac{1}{a^2} \times \sin ax + \sin ax \times \frac{1}{a^2} \times \left(\log \tan \frac{ax}{2} + \cos ax\right)$

$$= \frac{1}{a^2} \times \log \tan\left(\frac{ax}{2}\right)$$

Hence, the general solution of (1) is y = C.F+P.I.

i.e., $y = C_1 \cos ax + C_2 \sin ax + \frac{1}{a^2} \times \log \tan\left(\frac{ax}{2}\right)$.

(ii) Given, $\left(D^2 - 2D + 1\right)y = xe^x \log x, x > 0$, $D = \frac{d}{dx}$

Auxiliary equation is $(m-1)^2 = 0$

or $m = 1,1$

C.F $= (C_1 + C_2 x)e^x = C_1 e^x + C_2 x e^x$

Let $y_1 = e^x$, $y_2 = xe^x$. Also, $X = xe^x \log x$

Here, $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} e^x & xe^x \\ e^x & e^x + xe^x \end{vmatrix} = e^{2x} \neq 0$

Then, P. I $= u.y_1 + v.y_2$

where $u = -\int \frac{y_2.X}{W} dx = -\int \frac{xe^x.xe^x \log x}{e^{2x}} dx = -\int x^2 \log x dx$

$$= -\left[\log x.\frac{x^3}{3} - \int \frac{1}{x}.\frac{x^3}{3} dx\right]$$

$$= -\left[\log x.\frac{x^3}{3} - \frac{1}{9}x^3\right]$$

and $v = \int \frac{y_1.X}{W}dx = \int \frac{e^x.xe^x \log x}{e^{2x}}dx = \int x \log x dx$

$$= \frac{x^2}{2}\log x - \int \frac{x^2}{2}.\frac{1}{x}dx$$

$$= \frac{x^2}{2}\log x - \frac{x^2}{4}$$

P.I $= -e^x\left\{\frac{x^3}{3}\log x - \frac{x^3}{9}\right\} + xe^x\left\{\frac{x^2}{2}\log x - \frac{x^2}{4}\right\}$

$$= x^3e^x \log x\left(\frac{1}{2} - \frac{1}{3}\right) - x^3e^x\left(\frac{1}{4} - \frac{1}{9}\right)$$

$$= \left(\frac{1}{6}\right) \times x^3e^x \log x - \left(\frac{5}{36}\right) \times x^3e^x$$

Hence, the general solution is y = C.F+P.I.

$$\therefore\ y = C_1e^x + C_2xe^x + \left(\frac{1}{6}\right) \times x^3e^x \log x - \left(\frac{5}{36}\right) \times x^3e^x.$$

(iii) Do Yourself.

The required solution is

$$y = e^x\left(C_1 \cos x + C_2 \sin x\right) - e^x \cos x \log\left(\sin x + \tan x\right)$$

(iv) Do youself.

The required solution is

$$y = C_1e^{-x} + C_2e^{-2x} + \frac{x}{2} - \left(\frac{3}{4}\right) + \left(\frac{1}{10}\right) \times \left(3\sin x - \cos x\right).$$



12.8 MODEL QUESTIONS

Q 1 : Solve the following differential equations

(a) $(3x+2)^2\frac{d^2y}{dx^2} + 5(3x+2)\frac{dy}{dx} - 3y = x^2 + x + 1$

(b) $x^3\frac{d^3y}{dx^3}+3x^2\frac{d^2y}{dx^2}+x\frac{dy}{dx}+y=\log x+x$

(c) $\left(x^3D^3-3x^2D^2+6xD-6\right)y=\left(\log x\right)^2$

(d) $(x+1)^2\frac{d^2y}{dx^2}-3(x+1)+4y=x$

(e) $\left\{(1+2x)^2D^2-6(1+2x)D+16\right\}y=8(1+2x)^2$

Q 2 : Solve the following differential equations by the method of variation of parameters :

(a) $\frac{d^2y}{dx^2}-2\frac{dy}{dx}+y=2x$

(b) $\frac{d^2y}{dx^2}+y=\cos ecx\cot x$

(c) $\frac{d^2y}{dx^2}-2\frac{dy}{dx}+3y=x+\sin x$

(d) $\left(D^2-9\right)y=e^{2x}+x$

(e) $\frac{d^2y}{dx^2}+y=\sin^2 x$

*** ***** ***